

ON THE H -FUNCTION

Anatoly A. Kilbas

*Department of Mathematics and Mechanics, Belarusian State University
Minsk 220050, Belarus*

Megumi Saigo

*Department of Applied Mathematics, Fukuoka University
Fukuoka 814-0180, Japan*

Abstract

The paper is devoted to study the H -function defined by the Mellin-Barnes integral

$$H_{p,q}^{m,n}(z) = \frac{1}{2\pi i} \int_{\mathfrak{L}} \mathcal{H}_{p,q}^{m,n}(s) z^{-s} ds,$$

where the function $\mathcal{H}_{p,q}^{m,n}(s)$ is a certain ratio of products of Gamma functions with the argument s and the contour $\mathfrak{L}$ is specially chosen. The conditions for the existence of $H_{p,q}^{m,n}(z)$ are discussed and explicit power and power-logarithmic series expansions of $H_{p,q}^{m,n}(z)$ near zero and infinity are given. The obtained results define more precisely the known results.

Key words: H -function

AMS(MOS) Subject Classification: 33C40

1. Introduction

This paper deals with the H -function $H_{p,q}^{m,n}(z)$ introduced by Pincherle in 1888 (see [3, Section 1.19]). Interest in this function appeared in 1961, when Fox [4] investigated such a function as symmetrical Fourier kernel. Therefore, the H -function is often called Fox's H -function. For integers m, n, p, q such that

$0 \leq m \leq q, 0 \leq n \leq p$ and $a_i, b_j \in \mathbb{C}$ and $\alpha_i, \beta_j \in \mathbb{R}_+ = (0, \infty)$ ($1 \leq i \leq p, 1 \leq j \leq q$) the function is defined by the Mellin-Barnes integral

$$(1.1) \quad H_{p,q}^{m,n}(z) = \frac{1}{2\pi i} \int_{\mathfrak{L}} \mathcal{H}_{p,q}^{m,n}(s) z^{-s} ds,$$

where

$$(1.2) \quad \mathcal{H}_{p,q}^{m,n}(s) = \frac{\prod_{j=1}^m \Gamma(b_j + \beta_j s) \prod_{i=1}^n \Gamma(1 - a_i - \alpha_i s)}{\prod_{i=n+1}^p \Gamma(a_i + \alpha_i s) \prod_{j=m+1}^q \Gamma(1 - b_j - \beta_j s)},$$

the contour $\mathfrak{L}$ is specially chosen and an empty product, if it occurs, is taken to be one. The theory of this function may be found in [10], [2], [1], [9, Chapter 2], [12, Chapter 1] and [11, Section 8.3]. We only indicate that most of the elementary and special functions are particular cases of the H -function $H_{p,q}^{m,n}(z)$. In particular, if α 's and β 's are equal to 1, the H -function (1.1) reduces to Meijer's G -function $G_{p,q}^{m,n}(z)$.

The conditions of the existence of the H -function can be made by inspecting the convergence of the integral (1.1), which depend on the selection of the contour $\mathfrak{L}$ and the relations between parameters a_i, α_i ($i = 1, \dots, p$) and b_j, β_j ($j = 1, \dots, q$). Especially, the relations may depend on the numbers Δ, δ and μ defined by

$$(1.3) \quad \Delta = \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i,$$

$$(1.4) \quad \delta = \prod_{i=1}^p \alpha_i^{-\alpha_i} \prod_{j=1}^q \beta_j^{\beta_j}.$$

$$(1.5) \quad \mu = \sum_{j=1}^q b_j - \sum_{i=1}^p a_i + \frac{p-q}{2}.$$

Such a selection of $\mathfrak{L}$ and the relations on parameters are indicated in the handbook [11, Section 8.3.1], but some of the results there needs correction. In this paper we would like to give such a correction in the following cases:

(a) $\Delta \geq 0$ and the contour $\mathfrak{L} = \mathfrak{L}_{-\infty}$ in (1.1) runs from $-\infty + i\varphi_1$ to $-\infty + i\varphi_2$, $\varphi_1 < \varphi_2$, such that the poles of $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) lie on the left of $\mathfrak{L}_{-\infty}$ and those of $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) on the right of $\mathfrak{L}_{-\infty}$.

(b) $\Delta \leq 0$ and the contour $\mathfrak{L} = \mathfrak{L}_{+\infty}$ in (1.1) runs from $+\infty + i\varphi_1$ to $+\infty + i\varphi_2$, $\varphi_1 < \varphi_2$, such that the poles of $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) lie on the left of $\mathfrak{L}_{+\infty}$ and those of $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) on the right of $\mathfrak{L}_{+\infty}$.

Our results are based on the asymptotic behavior of the function $\mathcal{H}_{p,q}^{m,n}(s)$ given in (1.2) at infinity. Using the behavior and following [1], we give the series representation of $H_{p,q}^{m,n}(z)$ via residues of the integrand $\mathcal{H}_{p,q}^{m,n}(s)z^{-s}$. In this way we simplify the proof of Theorem 1 in [1] by applying the former results to find the explicit series expansions of $H_{p,q}^{m,n}(z)$. Such power expansions, as corollaries of the results from [1], were indicated in [12, Chapter 2.2] (see also [11, Section 8.3.1]), provided that the poles of Gamma-functions $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) and $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) do not coincide

$$(1.6) \quad \beta_j(a_i - 1 - k) \neq \alpha_i(b_j + l) \quad (i = 1, \dots, n; j = 1, \dots, m; k, l \in \mathbb{N}_0 = \{0, 1, 2, \dots\})$$

in the cases:

(c) $\Delta > 0$ with $z \neq 0$ or $\Delta = 0$ with $0 < |z| < \delta$, and the poles of Gamma-functions $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) are simple:

$$(1.7) \quad \beta_j(b_i + k) \neq \beta_i(b_j + l), \quad (i \neq j, \ i, j = 1, \dots, m; \ k, l \in \mathbb{N}_0);$$

(d) $\Delta < 0$ with $z \neq 0$ or $\Delta = 0$ with $|z| > \delta$, and the poles of Gamma-functions $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) are simple:

$$(1.8) \quad \alpha_j(1 - a_i + k) \neq \alpha_i(1 - a_j + l), \quad (i \neq j, \ i, j = 1, \dots, n; \ k, l \in \mathbb{N}_0).$$

When the poles of Gamma-functions in (c) and (d) coincide, explicit series expansions of $H_{p,q}^{m,n}(z)$ should be more complicated power-logarithmic expansions. Such expansions in particular cases of the Meijer's G -functions $G_{0,p}^{p,0}$ and $G_{p,p}^{p,0}$ and of the H -functions $H_{0,p}^{p,0}$ and $H_{p,p}^{p,0}$ were given in [7] and [8], respectively.

We obtain the explicit expansions of the H -function of general form $H_{p,q}^{m,n}(z)$ under the conditions in (1.6). We show that, if the poles of the Gamma-functions $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) and $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) coincide in the cases (c) and (d), respectively, then the H -function (1.1) has power-logarithmic series expansions. In particular, we give the asymptotic expansions of $H_{p,q}^{m,n}(z)$ near zero. We note that the obtained results will be different in the cases when either $\Delta \geq 0$ or $\Delta \leq 0$.

The paper is organized as follows. Section 2 is devoted to the conditions of the existence of the H -function (1.1) which are based on the asymptotic behavior of $\mathcal{H}_{p,q}^{m,n}(s)$ at infinity. Here we also give the representations of (1.1) via the residues of the integrand. The latter result is applied in Sections 3 and 4 to obtain the explicit power and power-logarithmic series expansions of $H_{p,q}^{m,n}(z)$ and, in particular, its asymptotic estimates near zero.

2. Existence and Representations of $H_{p,q}^{m,n}(z)$

First we give the asymptotic estimate of Gamma function $\Gamma(z)$, $z = x + iy$, [3, Chapter 1] at infinity on lines parallel to the coordinate axes.

Lemma 1. *Let $z = x + iy$ with $x, y \in \mathbb{R} = (-\infty, \infty)$. Then the following asymptotic estimates at infinity are valid:*

$$(2.1) \quad |\Gamma(x + iy)| \sim \sqrt{2\pi} |x|^{x-1/2} e^{-x-\pi(1-\text{sign } x)y/2} \quad (|x| \rightarrow \infty; \ y \neq 0 \text{ if } x < 0)$$

and

$$(2.2) \quad |\Gamma(x + iy)| \sim \sqrt{2\pi} |y|^{x-1/2} e^{-x-\pi|y|/2} \quad (|y| \rightarrow \infty).$$

Proof. Applying the Stirling formula [3, 1.18(2)]

$$(2.3) \quad \Gamma(z) \sim \sqrt{2\pi} e^{(z-1/2) \log z} e^{-z} \quad (|z| \rightarrow \infty; \ |\arg(z)| < \pi),$$

we have

$$(2.4) \quad |\Gamma(x + iy)| \sim \sqrt{2\pi} \left| e^{(x+iy-1/2)[\log|x+iy|+i\arg(x+iy)]} e^{-(x+iy)} \right| \\ \sim \sqrt{2\pi} |x + iy|^{x-1/2} e^{-x-y\arg(x+iy)} \quad (|x + iy| \rightarrow \infty; y \neq 0 \text{ if } x \leq 0).$$

Let $y \in \mathbb{R}$ be fixed and $|x| \rightarrow \infty$. Then $|x + iy| \sim |x|$, and $\arg(x + iy) \rightarrow 0$ as $x \rightarrow +\infty$ and $\arg(x + iy) \rightarrow \pi$ as $x \rightarrow -\infty$. Therefore, (2.4) implies

$$(2.5) \quad |\Gamma(x + iy)| \sim \sqrt{2\pi} |x|^{x-1/2} e^{-x} \quad (x \rightarrow +\infty)$$

and

$$(2.6) \quad |\Gamma(x + iy)| \sim \sqrt{2\pi} |x|^{x-1/2} e^{-x-\pi y} \quad (x \rightarrow -\infty; y \neq 0),$$

which yield (2.1).

Turning to the case $x \in \mathbb{R}$ being fixed and $|y| \rightarrow \infty$, we find $|x + iy| \sim |y|$ and $\arg(x + iy) \rightarrow \pi/2$ as $y \rightarrow \infty$ and $\arg(x + iy) \rightarrow -\pi/2$ as $y \rightarrow -\infty$. Thus (2.4) implies (2.2).

Remark 1. The relation [3, (1.18.6)] needs correction with addition of the multiplier e^x in the left hand side and it must be replaced by

$$(2.7) \quad \lim_{|y| \rightarrow \infty} |\Gamma(x + iy)| e^{x+\pi|y|/2} |y|^{1/2-x} = \sqrt{2\pi}.$$

Next assertion gives the asymptotic behavior of $\mathcal{H}_{p,q}^{m,n}(s)$ defined in (1.2) at infinity on lines parallel to the real axis.

Lemma 2. Let Δ, δ and μ be given by (1.3) to (1.5) and let $t, \sigma \in \mathbb{R}$. Then there hold the estimates

$$(2.8) \quad |\mathcal{H}_{p,q}^{m,n}(t + i\sigma)| \sim A \left(\frac{e}{t} \right)^{-\Delta t} \delta^t t^{\operatorname{Re}(\mu)} \quad (t \rightarrow +\infty)$$

with

$$(2.9) \quad A = (2\pi)^{m+n-(p+q)/2} e^{q-m-n} \frac{\prod_{j=1}^q [(\beta_j)^{\operatorname{Re}(b_j)-1/2} e^{-\operatorname{Re}(b_j)}]}{\prod_{i=1}^p [(\alpha_i)^{\operatorname{Re}(a_i)-1/2} e^{-\operatorname{Re}(a_i)}]} \frac{\prod_{i=1}^n e^{\pi[\sigma\alpha_i + \operatorname{Im}(a_i)]}}{\prod_{j=m+1}^q e^{\pi[\sigma\beta_j + \operatorname{Im}(b_j)]}},$$

and

$$(2.10) \quad |\mathcal{H}_{p,q}^{m,n}(t + i\sigma)| \sim B \left(\frac{e}{|t|} \right)^{\Delta|t|} \delta^{-|t|} |t|^{\operatorname{Re}(\mu)} \quad (t \rightarrow -\infty)$$

with

$$(2.11) \quad B = (2\pi)^{m+n-(p+q)/2} e^{q-m-n} \frac{\prod_{j=1}^q [(\beta_j)^{\operatorname{Re}(b_j)-1/2} e^{-\operatorname{Re}(b_j)}]}{\prod_{i=1}^p [(\alpha_i)^{\operatorname{Re}(a_i)-1/2} e^{-\operatorname{Re}(a_i)}]} \frac{\prod_{i=n+1}^p e^{\pi[\sigma\alpha_i + \operatorname{Im}(a_i)]}}{\prod_{j=1}^m e^{\pi[\sigma\beta_j + \operatorname{Im}(b_j)]}}.$$

Proof. By virtue of (2.1), we have, for a variable $s = t + i\sigma$ and a complex constant $k = c + id$,

$$(2.12) \quad |\Gamma(s + k)| \sim \sqrt{2\pi} t^{t+c-1/2} e^{-(t+c)} \quad (t \rightarrow +\infty)$$

and

$$(2.13) \quad |\Gamma(s + k)| \sim \sqrt{2\pi} |t|^{t+c-1/2} e^{-(t+c)} e^{-\pi(\sigma+d)} \quad (t \rightarrow -\infty).$$

Substituting these estimates into (1.2) and using (1.3) to (1.5), we obtain (2.8) and (2.10).

Remark 2. The asymptotic estimate of the function $\mathcal{H}_{p,q}^{m,n}(s)$ at infinity on lines parallel to the imaginary axis $\mathcal{H}_{p,q}^{m,n}(\sigma + it)$ as $|t| \rightarrow \infty$ was given in our paper with Shlapakov [5].

By appealing to Lemma 2, we give conditions of the existence of the H -function (1.1) with the contour $\mathfrak{L}$ being chosen as indicated in (a) and (b) in Section 1.

Theorem 1. Let Δ, δ and μ be given by (1.3) to (1.5). Then the function $H_{p,q}^{m,n}(z)$ defined by (1.1) and (1.2) exists in the following cases:

$$(2.14) \quad \mathfrak{L} = \mathfrak{L}_{-\infty}, \quad \Delta > 0, \quad z \neq 0;$$

$$(2.15) \quad \mathfrak{L} = \mathfrak{L}_{-\infty}, \quad \Delta = 0, \quad 0 < |z| < \delta; \text{ label 1.2.15}$$

$$(2.16) \quad \mathfrak{L} = \mathfrak{L}_{-\infty}, \quad \Delta = 0, \quad |z| = \delta, \quad \operatorname{Re}(\mu) < -1;$$

$$(2.17) \quad \mathfrak{L} = \mathfrak{L}_{+\infty}, \quad \Delta < 0, \quad z \neq 0;$$

$$(2.18) \quad \mathfrak{L} = \mathfrak{L}_{+\infty}, \quad \Delta = 0, \quad |z| > \delta;$$

$$(2.19) \quad \mathfrak{L} = \mathfrak{L}_{+\infty}, \quad \Delta = 0, \quad |z| = \delta, \quad \operatorname{Re}(\mu) < -1.$$

Proof. Let us first consider the case (a) for which $\Delta \geq 0$ and $\mathfrak{L} = \mathfrak{L}_{-\infty}$. We have to investigate the convergence of the integral (1.1) on the lines

$$(2.20) \quad l_1 = \{t \in \mathbb{R} : t + i\varphi_1\} \quad \text{and} \quad l_2 = \{t \in \mathbb{R} : t + i\varphi_2\} \quad \text{for} \quad \varphi_1 < \varphi_2$$

as $t \rightarrow -\infty$. According to (2.10), we have the following asymptotic estimate for the integrand of (1.1):

$$(2.21) \quad |\mathcal{H}_{p,q}^{m,n}(s) z^{-s}| \sim B_i e^{\varphi_i \arg z} \left(\frac{e}{|t|} \right)^{\Delta|t|} \left(\frac{|z|}{\delta} \right)^{|t|} |t|^{\operatorname{Re}(\mu)} \quad (t \rightarrow -\infty; t \in l_i \ (i = 1, 2)),$$

where B_1 and B_2 are given by (2.11) with σ being replaced by φ_1 and φ_2 , respectively. It follows from (2.21) that the integral (1.1) is convergent if and only if one of the conditions in (2.14) to (2.16) is satisfied.

In the case (b), $\Delta \leq 0$ and the contour $\mathfrak{L}$ is taken to be $\mathfrak{L}_{+\infty}$. Then we have to investigate the convergence of the integral (1.1) on the lines l_1 and l_2 in (2.20), as $t \rightarrow +\infty$. By virtue of (2.8) and (2.9), we have the asymptotic estimate:

$$(2.22) \quad |\mathcal{H}_{p,q}^{m,n}(s)z^{-s}| \sim A_i e^{\varphi_i \arg z} \left(\frac{e}{t}\right)^{-\Delta t} \left(\frac{\delta}{|z|}\right)^t t^{\operatorname{Re}(\mu)} \quad (t \rightarrow +\infty; t \in l_i \ (i = 1, 2)),$$

where A_1 and A_2 are given by (2.9) with σ being replaced by φ_1 and φ_2 , respectively. Thus (2.22) implies that the integral (1.1) converges if and only if one of the conditions in (2.17) to (2.19) holds.

Corollary 1. *The estimate (2.21) holds for $t \rightarrow -\infty$ uniformly on sets which have a positive distance to the points*

$$(2.23) \quad b_{jl} = -\frac{b_j + l}{\beta_j} \quad (j = 1, \dots, m; l \in \mathbb{N}_0).$$

and do not contain points to the right of $\mathfrak{L}_{-\infty}$.

The estimate (2.22) holds for $t \rightarrow +\infty$ uniformly on sets which have a positive distance to the points

$$(2.24) \quad a_{ik} = \frac{1 - a_i + k}{\alpha_i} \quad (i = 1, \dots, n; k \in \mathbb{N}_0).$$

and do not contain points to the left of $\mathfrak{L}_{+\infty}$.

Remark 3. The conditions for the existence of the H -function (1.1)

$$(2.25) \quad \sum_{i=1}^n \alpha_i - \sum_{i=n+1}^p \alpha_i + \sum_{j=1}^m \beta_j - \sum_{j=m+1}^q \beta_j \geq 0, \quad \operatorname{Re}(\mu) < 0$$

given in [11, Section 8.3.1] in the cases when $\mathfrak{L} = \mathfrak{L}_{-\infty}, \Delta = 0, |z| = \delta$ ([8, Section 8.3.1.3]) and $\mathfrak{L} = \mathfrak{L}_{+\infty}, \Delta = 0, |z| = \delta$ ([8, Section 8.3.1.4]) can be replaced by the condition

$$(2.26) \quad \operatorname{Re}(\mu) < -1.$$

The following statement follows from Theorem 1, Corollary 1 and the theory of residues.

Theorem 2. (A) *If the conditions in (1.6), and (2.14) or (2.15) are satisfied, then the H -function (1.1) is an analytic function of z in the corresponding domain indicated in (2.14) or (2.15), and*

$$(2.27) \quad H_{p,q}^{m,n}(z) = \sum_{j=1}^m \sum_{l=0}^{\infty} \operatorname{Res}_{s=b_{jl}} [\mathcal{H}_{p,q}^{m,n}(s)z^{-s}],$$

where b_{jl} are given in (2.23).

(B) If the conditions in (1.6), and (2.17) or (2.18) are satisfied, then the H -function (1.1) is an analytic function of z in the corresponding domain indicated in (2.17) or (2.18), and

$$(2.28) \quad H_{p,q}^{m,n}(z) = - \sum_{i=1}^n \sum_{k=0}^{\infty} \text{Res}_{s=a_{ik}} [\mathcal{H}_{p,q}^{m,n}(s) z^{-s}],$$

where a_{ik} are given in (2.24).

Remark 4. The first assertion of Theorem 2 was proved in [1, p.278, Theorem 1] for the H -function represented by the integral obtained from (1.1) and (1.2) after replacing s by $-s$. The proof of Theorem 1 in [1] is complicated and based on Lemma 2 there in which the asymptotic estimate at infinity of the functions $h_0(s)$ defined by

$$(2.29) \quad h_0(s) = \frac{\prod_{i=1}^p \Gamma(1 - a_i + \alpha_i s)}{\prod_{j=1}^q \Gamma(1 - b_j + \beta_j s)}$$

is given. But our proof of Theorem 2 along the ideas of [1] is more simple and is based on the asymptotic estimate of $\mathcal{H}_{p,q}^{m,n}(s)$ at infinity given in Lemma 2.

3. Explicit Power Series Expansions

In this section we apply Theorem 2 to obtain explicit power series expansions of the H -function (1.1) under the condition (1.6) in the case of (1.7) or (1.8).

First we consider the former case. By Theorem 2(A), we have to evaluate the residues of $\mathcal{H}(s)z^{-s}$ at the points $s = b_{jl}$ given in (2.23), where and in what follows we simplify $\mathcal{H}_{p,q}^{m,n}(s)$ by $\mathcal{H}(s)$. To evaluate these residues we use the property of the Gamma-function [6, (3.30)], that is, in a neighbourhood of the poles $z = -k$ ($k \in \mathbb{N}_0$) the Gamma-function $\Gamma(z)$ can be expanded in powers of $z + k = \epsilon$

$$(3.1) \quad \Gamma(z) = \frac{(-1)^k}{k! \epsilon} [1 + \epsilon \psi(1+k) + O(\epsilon^2)], \quad \text{where} \quad \psi(z) = \frac{\Gamma'(z)}{\Gamma(z)}.$$

Since the poles b_{jl} are simple, i.e., the conditions in (1.7) hold,

$$(3.2) \quad \text{Res}_{s=b_{jl}} [\mathcal{H}(s)z^{-s}] = h_{jl}^* z^{-b_{jl}} \quad (j = 1, \dots, m; \quad l \in \mathbb{N}_0),$$

where

$$(3.3) \quad h_{jl}^* = \lim_{s \rightarrow b_{jl}} [(s - b_{jl}) \mathcal{H}(s)] \\ = \frac{(-1)^l}{l! \beta_j} \frac{\prod_{i=1, i \neq j}^m \Gamma\left(b_i - [b_j + l] \frac{\beta_i}{\beta_j}\right) \prod_{i=1}^n \Gamma\left(1 - a_i + [b_j + l] \frac{\alpha_i}{\beta_j}\right)}{\prod_{i=n+1}^p \Gamma\left(a_i - [b_j + l] \frac{\alpha_i}{\beta_j}\right) \prod_{i=m+1}^q \Gamma\left(1 - b_i + [b_j + l] \frac{\beta_i}{\beta_j}\right)}.$$

Thus we obtain

Theorem 3. *Let the conditions in (1.6) and (1.7) be satisfied and let either $\Delta > 0, z \neq 0$ or $\Delta = 0, 0 < |z| < \delta$. Then the H -function (1.1) has the power series expansion*

$$(3.4) \quad H_{p,q}^{m,n}(z) = \sum_{j=1}^m \sum_{l=0}^{\infty} h_{jl}^* z^{(b_j+l)/\beta_j},$$

where the constants h_{jl}^* are given by (3.3).

Corollary 2. *If the conditions in (1.6) and (1.7) are satisfied and $\Delta \geq 0$, then (3.4) gives the asymptotic expansion of $H_{p,q}^{m,n}(z)$ near zero and the main terms of this asymptotic formula have the form:*

$$(3.5) \quad H_{p,q}^{m,n}(z) = \sum_{j=1}^m \left[h_j^* z^{b_j/\beta_j} + O\left(z^{(b_j+1)/\beta_j}\right) \right] \quad (z \rightarrow 0),$$

where

$$(3.6) \quad h_j^* \equiv h_{j0}^* = \frac{1}{\beta_j} \frac{\prod_{i=1, i \neq j}^m \Gamma\left(b_i - \frac{b_j \beta_i}{\beta_j}\right) \prod_{i=1}^n \Gamma\left(1 - a_i + \frac{b_j \alpha_i}{\beta_j}\right)}{\prod_{i=n+1}^p \Gamma\left(a_i - \frac{b_j \alpha_i}{\beta_j}\right) \prod_{i=m+1}^q \Gamma\left(1 - b_i + \frac{b_j \beta_i}{\beta_j}\right)}.$$

Corollary 3. *Let the conditions in (1.6) and (1.7) be satisfied, and let $\Delta \geq 0$ and j_0 ($1 \leq j_0 \leq m$) be an integer such that*

$$(3.7) \quad \frac{\operatorname{Re}(b_{j_0})}{\beta_{j_0}} = \min_{1 \leq j \leq m} \left[\frac{\operatorname{Re}(b_j)}{\beta_j} \right].$$

Then there holds the asymptotic estimate:

$$(3.8) \quad H_{p,q}^{m,n}(z) = h_{j_0}^* z^{b_{j_0}/\beta_{j_0}} + o\left(z^{b_{j_0}/\beta_{j_0}}\right) \quad (z \rightarrow 0),$$

where $h_{j_0}^*$ is given by (3.6) with $j = j_0$. In particular,

$$(3.9) \quad H_{p,q}^{m,n}(z) = O(z^\rho) \quad (z \rightarrow 0) \quad \text{with} \quad \rho = \min_{1 \leq j \leq m} \left[\frac{\operatorname{Re}(b_j)}{\beta_j} \right].$$

Now we consider the case (1.8) when the poles of the Gamma-functions $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) are simple. By (3.1), evaluating the residues of $\mathcal{H}(s)z^{-s}$ at the points a_{ik} given in (2.24) we have similarly to the previous argument that

$$(3.10) \quad \operatorname{Res}_{s=a_{ik}} [\mathcal{H}(s)z^{-s}] = -h_{ik} z^{-a_{ik}} \quad (i = 1, \dots, n; k \in \mathbb{N}_0),$$

where a_{ik} are given by (2.24) and

$$(3.11) \quad h_{ik} = \lim_{s \rightarrow a_{ik}} [-(s - a_{ik})\mathcal{H}(s)] \\ = \frac{(-1)^k}{k! \alpha_i} \frac{\prod_{j=1}^m \Gamma\left(b_j + [1 - a_i + k] \frac{\beta_j}{\alpha_i}\right) \prod_{j=1, j \neq i}^n \Gamma\left(1 - a_j - [1 - a_i + k] \frac{\alpha_j}{\alpha_i}\right)}{\prod_{j=n+1}^p \Gamma\left(a_j + [1 - a_i + k] \frac{lpha_j}{\alpha_i}\right) \prod_{j=m+1}^q \Gamma\left(1 - b_j - [1 - a_i + k] \frac{\beta_j}{\alpha_i}\right)}.$$

Thus from Theorem 2(B) we have

Theorem 4. *Let the conditions in (1.6) and (1.8) be satisfied and let either $\Delta < 0, z \neq 0$ or $\Delta = 0, |z| > \delta$. Then the H -function (1.1) has the power series expansion*

$$(3.12) \quad H_{p,q}^{m,n}(z) = \sum_{i=1}^n \sum_{k=0}^{\infty} h_{ik} z^{(a_i - k - 1)/\alpha_i},$$

where the constants h_{ik} are given by (3.11).

Corollary 4. *If the conditions in (1.6) and (1.8) are satisfied and $\Delta < 0$, then (3.12) gives the asymptotic expansion of $H_{p,q}^{m,n}(z)$ near infinity and the main terms of this asymptotic formula have the form:*

$$(3.13) \quad H_{p,q}^{m,n}(z) = \sum_{i=1}^n \left[h_i z^{(a_i - 1)/\alpha_i} + O\left(z^{(a_i - 2)/\alpha_i}\right) \right] \quad (|z| \rightarrow \infty),$$

where

$$(3.14) \quad h_i \equiv h_{i0} = \frac{1}{\alpha_i} \frac{\prod_{j=1}^m \Gamma\left(b_j - [a_i - 1] \frac{\beta_j}{\alpha_i}\right) \prod_{j=1, j \neq i}^n \Gamma\left(1 - a_j + [a_i - 1] \frac{\alpha_j}{\alpha_i}\right)}{\prod_{j=n+1}^p \Gamma\left(a_j - [a_i - 1] \frac{\alpha_j}{\alpha_i}\right) \prod_{j=m+1}^q \Gamma\left(1 - b_j + [a_i - 1] \frac{\beta_j}{\alpha_i}\right)}.$$

Corollary 5. *Let the conditions in (1.6) and (1.8) be satisfied, and let $\Delta < 0$ and i_0 ($1 \leq i_0 \leq n$) be an integer such that*

$$(3.15) \quad \frac{\operatorname{Re}(a_{i_0}) - 1}{\alpha_{i_0}} = \max_{1 \leq i \leq n} \left[\frac{\operatorname{Re}(a_i) - 1}{\alpha_i} \right].$$

Then there holds the asymptotic estimate:

$$(3.16) \quad H_{p,q}^{m,n}(z) = h_{i_0} z^{(a_{i_0} - 1)/\alpha_{i_0}} + o\left(z^{(a_{i_0} - 1)/\alpha_{i_0}}\right) \quad (|z| \rightarrow \infty),$$

where h_{i_0} is given by (3.14) with $i = i_0$. In particular,

$$(3.17) \quad H_{p,q}^{m,n}(z) = O(z^\varrho) \quad (|z| \rightarrow \infty) \quad \text{with} \quad \varrho = \max_{1 \leq i \leq n} \left[\frac{\operatorname{Re}(a_i) - 1}{\alpha_i} \right].$$

Remark 5. The relations (3.4) and (3.12) are given in [12, (2.2.4) and (2.2.7)] and [11, 8.3.2.3 and 8.3.2.4].

4. Explicit Power-Logarithmic Series Expansions

Now let us discuss the case when the condition (1.6) holds, but (1.7) or (1.8) is violated:

(e) $\mathfrak{L} = \mathfrak{L}_{-\infty}, \Delta \geq 0$ and some poles of the

Gamma-functions $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) coincide.

(f) $\mathfrak{L} = \mathfrak{L}_{+\infty}, \Delta \leq 0$ and some poles of the Gamma-functions $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) coincide.

First we consider the case (e). Let $b \equiv b_{jl}$ be one of points (2.23) for which some poles of the Gamma-functions $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) coincide and $N^* \equiv N_{jl}^*$ be order of this pole. It means that there exist $j_1, \dots, j_{N^*} \in \{1, \dots, m\}$ and $l_{j_1}, \dots, l_{j_{N^*}} \in \mathbb{N}_0$ such that

$$(4.1) \quad b = b_{jl} = -\frac{b_{j_1} + l_{j_1}}{\beta_{j_1}} = \dots = -\frac{b_{j_{N^*}} + l_{j_{N^*}}}{\beta_{j_{N^*}}}.$$

Then $\mathcal{H}(s)z^{-s}$ has the pole of order N^* at b and hence

$$(4.2) \quad \text{Res}_{s=b}[\mathcal{H}(s)z^{-s}] = \frac{1}{(N^* - 1)!} \lim_{s \rightarrow b} [(s - b)^{N^*} \mathcal{H}(s)z^{-s}]^{(N^* - 1)}.$$

We denote

$$(4.3) \quad \mathcal{H}_1^*(s) = (s - b)^{N^*} \prod_{j=j_1}^{j_{N^*}} \Gamma(b_j + \beta_j s), \quad \mathcal{H}_2^*(s) = \frac{\mathcal{H}(s)}{\prod_{j=j_1}^{j_{N^*}} \Gamma(b_j + \beta_j s)}.$$

Using the Leibniz rule, we have

$$\begin{aligned} [(s - b)^{N^*} \mathcal{H}(s)z^{-s}]^{(N^* - 1)} &= \sum_{n=0}^{N^* - 1} N^* - 1 \text{ choosen} [\mathcal{H}_1^*(s)]^{(N^* - 1 - n)} [\mathcal{H}_2^*(s)z^{-s}]^{(n)} \\ &= \sum_{n=0}^{N^* - 1} \binom{N^* - 1}{n} [\mathcal{H}_1^*(s)]^{(N^* - 1 - n)} \sum_{i=0}^n \binom{n}{i} (-1)^i [\mathcal{H}_2^*(s)]^{(n - i)} z^{-s} [\log z]^i \\ &= z^{-s} \sum_{i=0}^{N^* - 1} \left\{ \sum_{n=i}^{N^* - 1} (-1)^i \binom{N^* - 1}{n} \binom{n}{i} [\mathcal{H}_1^*(s)]^{(N^* - 1 - n)} [\mathcal{H}_2^*(s)]^{(n - i)} \right\} [\log z]^i. \end{aligned}$$

Substituting this into (4.2), we obtain

$$(4.4) \quad \text{Res}_{s=b_{jl}}[\mathcal{H}(s)z^{-s}] = z^{(b_j + l)/\beta_j} \sum_{i=0}^{N_{jl}^* - 1} H_{jli}^* [\log z]^i,$$

where

$$(4.5) \quad H_{jli}^* \equiv H_{jli}^*(N_{jl}^*; b_{jl}) \\ = \frac{1}{(N_{jl}^* - 1)!} \sum_{n=i}^{N_{jl}^* - 1} (-1)^i \binom{N_{jl}^* - 1}{n} \binom{n}{i} [\mathcal{H}_1^*(b_{jl})]^{(N_{jl}^* - 1 - n)} [\mathcal{H}_2^*(b_{jl})]^{(n-i)}.$$

In particular, if $l = 0$ and $i = N_{j0}^* - 1$, then by setting $N_{j0}^* \equiv N_j^*$ and from (3.1), (4.1) and (4.3), we have

$$(4.6) \quad H_j^* \equiv H_{j,0,N_j^*-1}^*(N_j^*; b_{j0}) = \frac{(-1)^{N_j^*-1}}{(N_j^* - 1)!} \mathcal{H}_1^*(b_{j0}) \mathcal{H}_2^*(b_{j0}) \\ = \frac{(-1)^{N_j^*-1}}{(N_j^* - 1)!} \left\{ \prod_{k=1}^{N_j^*} \frac{(-1)^{j_k}}{j_k! \beta_{j_k}} \right\} \frac{\prod_{i=1, i \neq j_1, \dots, j_{N_j^*}}^m \Gamma\left(b_i - \frac{b_j \beta_i}{\beta_j}\right) \prod_{i=1}^n \Gamma\left(1 - a_i + \frac{b_j \alpha_i}{\beta_j}\right)}{\prod_{i=n+1}^p \Gamma\left(a_i - \frac{b_j \alpha_i}{\beta_j}\right) \prod_{i=m+1}^q \Gamma\left(1 - b_i + \frac{b_j \beta_i}{\beta_j}\right)}.$$

Thus, in view of Theorem 2(A), we have

Theorem 5. Let the conditions in (1.6) be satisfied and let either $\Delta > 0, z \neq 0$ or $\Delta = 0, 0 < |z| < \delta$. Then the H -function (1.1) has the power-logarithmic series expansion

$$(4.7) \quad H_{p,q}^{m,n}(z) = \sum'_{j,l} h_{jl}^* z^{(b_j+l)/\beta_j} + \sum''_{j,l} \sum_{i=0}^{N_{jl}^*-1} H_{jli}^* z^{(b_j+l)/\beta_j} [\log z]^i.$$

Here $\sum'$ and $\sum''$ are summations taken over j, l ($j = 1, \dots, m; l = 0, 1, \dots$) such that the Gamma-functions $\Gamma(b_j + \beta_j s)$ have simple poles and poles of order N_{jl}^* at the points b_{jl} , respectively, and the constants h_{jl}^* are given by (3.3) while the constants H_{jli}^* are given by (4.5).

Corollary 6. If the conditions in (1.6) are satisfied and $\Delta \geq 0$, then (4.7) gives the asymptotic expansion of $H_{p,q}^{m,n}(z)$ near zero and the main terms of this asymptotic formula have the form:

$$(4.8) \quad H_{p,q}^{m,n}(z) = \sum'_{j=1}^m \left[h_j^* z^{b_j/\beta_j} + O(z^{(b_j+1)/\beta_j}) \right] \\ + \sum''_{j=1}^m \left(H_j^* z^{b_j/\beta_j} [\log z]^{N_j^*-1} + O(z^{(b_j+1)/\beta_j} [\log z]^{N_j^*-1}) \right) \quad (z \rightarrow 0).$$

Here $\sum'$ and $\sum''$ are summations taken over j ($j = 1, \dots, m$) such that the Gamma-functions $\Gamma(b_j + \beta_j s)$ have simple poles and poles of order $N_j^* \equiv N_{j0}^*$ at the points b_{j0} , respectively, and h_j^* are given by (3.6) while H_j^* are given by (4.6).

Corollary 7. *Let the conditions in (1.6) be satisfied, and let $\Delta \geq 0$ and b_{jl} be poles of the Gamma-functions $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$). Let j_{01} and j_{02} ($1 \leq j_{01}, j_{02} \leq m$) be integers such that*

$$(4.9) \quad \frac{\operatorname{Re}(b_{j_{01}})}{\beta_{j_{01}}} = \min_{1 \leq j \leq m} \left[\frac{\operatorname{Re}(b_j)}{\beta_j} \right],$$

when the poles b_{jl} ($j = 1, \dots, m; l \in \mathbb{N}_0$) are simple, and

$$(4.10) \quad \frac{\operatorname{Re}(b_{j_{02}})}{\beta_{j_{02}}} = \min_{1 \leq j \leq m} \left[\frac{\operatorname{Re}(b_j)}{\beta_j} \right],$$

when the poles b_{jl} ($j = 1, \dots, m; l \in \mathbb{N}_0$) coincide.

a) *If $j_{01} < j_{02}$, then the asymptotic expansion of the H -function has the form:*

$$(4.11) \quad H_{p,q}^{m,n}(z) = h_{j_{01}}^* z^{b_{j_{01}}/\beta_{j_{01}}} + o\left(z^{b_{j_{01}}/\beta_{j_{01}}}\right) \quad (z \rightarrow 0),$$

where $h_{j_0}^*$ is given by (3.6) with $j = j_{01}$. In particular, the relation (3.9) holds.

b) *If $j_{01} \geq j_{02}$ and $b_{j_{02},0}$ has the pole of order $N_{j_{02}}^*$, then the first term in asymptotic expansion of the H -function has the form:*

$$(4.12) \quad H_{p,q}^{m,n}(z) = H_{j_{02}}^* z^{b_{j_{02}}/\beta_{j_{02}}} [\log(z)]^{N_{j_{02}}^* - 1} + o\left(z^{b_{j_{02}}/\beta_{j_{02}}} [\log(z)]^{N_{j_{02}}^* - 1}\right) \quad (z \rightarrow 0),$$

where $H_{j_{02}}^*$ is given by (4.6) with $j = j_{02}$. In particular, if N^* is the largest order of general poles of Gamma-functions $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$), then

$$(4.13) \quad H_{p,q}^{m,n}(z) = O\left(z^\rho [\log(z)]^{N^*}\right) \quad (z \rightarrow 0) \quad \text{with} \quad \rho = \min_{1 \leq j \leq m} \left[\frac{\operatorname{Re}(b_j)}{\beta_j} \right].$$

Now we consider the case (f). Let $a = a_{ik}$ be one of points (2.24) for which some poles of $\Gamma(1 - a_i - \alpha_i s)$ coincide and $N = N_{ik}$ be order of this pole. It means that there exist $i_1, \dots, i_N \in \{1, \dots, n\}$ and $k_{i_1}, \dots, k_{i_N} \in \mathbb{N}_0$ such that

$$(4.14) \quad a = a_{ik} \equiv \frac{1 - a_{i_1} + k_{i_1}}{\alpha_{i_1}} = \dots = \frac{1 - a_{i_N} + k_{i_N}}{\alpha_{i_N}}.$$

Then the integrand $\mathcal{H}(s)z^{-s}$ of the integral (1.1) has the pole of order N at a . Similarly to (4.3), we denote

$$(4.15) \quad \mathcal{H}_1(s) = (s - a)^N \prod_{i=i_1}^{i_N} \Gamma(1 - a_i - \alpha_i s), \quad \mathcal{H}_2(s) = \frac{\mathcal{H}(s)}{\prod_{i=i_1}^{i_N} \Gamma(1 - a_i - \alpha_i s)}$$

and, then, find similarly to (4.4) and (4.5) that

$$(4.16) \quad \text{Res}_{s=a_{ik}} [\mathcal{H}(s)z^{-s}] = z^{(a_i-1-k)/\alpha_i} \sum_{j=0}^{N_{ik}-1} H_{ikj} [\log(z)]^j,$$

where

$$(4.17) \quad H_{ikj} \equiv H_{ikj}(N_{ik}; a_{ik}) \\ = \frac{1}{(N_{ik}-1)!} \left\{ \sum_{n=j}^{N_{ik}-1} (-1)^j \binom{N_{ik}-1}{n} \binom{n}{j} [\mathcal{H}_1(a_{ik})]^{(N_{ik}-1-n)} [\mathcal{H}_2(a_{ik})]^{(n-j)} \right\}.$$

In particular, if we set $k = 0$, $j = N_{i0} - 1$ and $N_{i0} \equiv N_i$, then, using (4.15) and (3.1), we have

$$(4.18) \quad H_i \equiv H_{i,0,N_i-1}(N_i; a_{i0}) = \frac{(-1)^{N_i-1}}{(N_i-1)!} \mathcal{H}_1(a_{i0}) \mathcal{H}_2(a_{i0}) = \frac{(-1)^{N_i-1}}{(N_i-1)!} \\ \times \left(\prod_{k=1}^{N_i} \frac{(-1)^{i_k-1}}{i_k! \alpha_{i_k}} \right) \frac{\prod_{j=1}^m \Gamma\left(b_j + [1 - a_i] \frac{\beta_j}{\alpha_i}\right) \prod_{j=1, j \neq i_1, \dots, i_{N_i}}^n \Gamma\left(1 - a_j - [1 - a_i] \frac{\alpha_j}{\alpha_i}\right)}{\prod_{j=n+1}^p \Gamma\left(a_j + [1 - a_i] \frac{\alpha_j}{\alpha_i}\right) \prod_{j=m+1}^q \Gamma\left(1 - b_j - [1 - a_i] \frac{\beta_j}{\alpha_i}\right)}.$$

Therefore, Theorem 2(B) implies the similar result to Theorem 5:

Theorem 6. *Let the conditions in (1.6) be satisfied and let either $\Delta < 0, z \neq 0$ or $\Delta = 0, |z| > \delta$. Then the H -function (1.1) has the power-logarithmic series expansion*

$$(4.19) \quad H_{p,q}^{m,n}(z) = \sum'_{i,k} h_{ik} z^{(a_i-1-k)/\alpha_i} + \sum''_{i,k} \sum_{j=0}^{N_{ik}-1} H_{ikj} z^{(a_i-1-k)/\alpha_i} [\log(z)]^j.$$

Here $\sum'$ and $\sum''$ are summations taken over i, k ($i = 1, \dots, n$; $k = 0, 1, \dots$) such that Gamma-functions $\Gamma(1 - a_i - \alpha_i s)$ have simple poles and poles of order N_{ik} at the points a_{ik} , respectively, and the constants h_{ik} are given by (3.11) while the constants H_{ikj} are given by (4.17).

Corollary 8. *If the conditions in (1.6) are satisfied and $\Delta \leq 0$, then (4.19) gives the asymptotic expansion of $H_{p,q}^{m,n}(z)$ near infinity and the main terms of this asymptotic formula have the form:*

$$(4.20) \quad H_{p,q}^{m,n}(z) = \sum_{i=1}^n \left[h_i z^{(a_i-1)/\alpha_i} + O(z^{(a_i-2)/\alpha_i}) \right] \\ + \sum_{i=1}^n \left(H_i z^{(a_i-1)/\alpha_i} [\log(z)]^{N_i-1} + O(z^{(a_i-2)/\alpha_i} [\log z]^{N_i-1}) \right) \quad (|z| \rightarrow \infty).$$

Here $\sum'$ and $\sum''$ are summations taken over i ($i = 1, \dots, n$) such that the Gamma-functions $\Gamma(1 - a_i - \alpha_i s)$ have simple poles and poles of order $N_i \equiv N_{i0}$ in the points a_{i0} in (2.24), respectively, and h_i are given by (3.14) while H_i are given by (4.18).

Corollary 9. *Let the conditions in (1.6) be satisfied, and let $\Delta \leq 0$ and a_{ik} be poles of the Gamma-function $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$). Let i_{01} and i_{02} ($1 \leq i_{01}, i_{02} \leq n$) be integers such that*

$$(4.21) \quad \frac{\operatorname{Re}(a_{i_{01}}) - 1}{\alpha_{i_{01}}} = \max_{1 \leq i \leq n} \left[\frac{\operatorname{Re}(a_i) - 1}{\alpha_i} \right],$$

when the poles a_{ik} ($i = 1, \dots, n; k = 0, 1, \dots$) are simple, and

$$(4.22) \quad \frac{\operatorname{Re}(a_{i_{02}}) - 1}{\alpha_{i_{02}}} = \max_{1 \leq i \leq n} \left[\frac{\operatorname{Re}(a_i) - 1}{\alpha_i} \right],$$

when the poles a_{ik} ($i = 1, \dots, n; k \in \mathbb{N}_0$) coincide.

(a) *If $i_{01} < i_{02}$, then the asymptotic expansion of the H -function has the form:*

$$(4.23) \quad H_{p,q}^{m,n}(z) = h_{i_{01}} z^{(a_{i_{01}} - 1)/\alpha_{i_{01}}} + o\left(z^{(a_{i_{01}} - 1)/\alpha_{i_{01}}}\right) \quad (|z| \rightarrow \inf ty),$$

where $h_{i_{01}}$ is given by (3.14) with $i = i_{01}$. In particular, the relation (3.17) holds.

(b) *If $i_{01} \geq i_{02}$ and $a_{i_{02},0}$ has the pole of order $N_{i_{02}}$, then the asymptotic expansion of the H -function has the form:*

$$(4.24) \quad H_{p,q}^{m,n}(z) = H_{i_{02}} z^{(a_{i_{02}} - 1)/\alpha_{i_{02}}} [\log z]^{N_{i_{02}} - 1} + o\left(z^{(a_{i_{02}} - 1)/\alpha_{i_{02}}} [\log(z)]^{N_{i_{02}} - 1}\right) \quad (|z| \rightarrow \infty),$$

where $H_{i_{02}}$ is given by (4.18) with $i = i_{02}$. In particular, if N is the smallest order of general poles of Gamma-functions $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$), then

$$(4.25) \quad H_{p,q}^{m,n}(z) = O\left(z^\varrho [\log(z)]^N\right) \quad (|z| \rightarrow \infty) \quad \text{with} \quad \varrho = \min_{1 \leq i \leq n} \left[\frac{\operatorname{Re}(a_i) - 1}{\alpha_i} \right].$$

In conclusion, we give the following consequence of Corollaries 3, 5, 7 and 9, which unifies the power and power-logarithmic asymptotic behavior of $H_{p,q}^{m,n}(z)$ near zero and infinity.

Theorem 7. *Let the conditions in (1.6) be satisfied.*

(a) *If $\Delta \geq 0$ and the poles of the Gamma-function $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) are simple, then the H -function (1.3) has the asymptotic estimate (3.9) at zero. If some of poles of $\Gamma(b_j + \beta_j s)$ ($j = 1, \dots, m$) coincide, then $H_{p,q}^{m,n}(z)$ has the asymptotic estimate either (3.9) or (4.13) at zero.*

(b) *If $\Delta \leq 0$ and the poles of the Gamma-function $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) are simple, then the H -function (1.3) has the asymptotic estimate (3.17) at infinity. If some of poles of $\Gamma(1 - a_i - \alpha_i s)$ ($i = 1, \dots, n$) coincide, then $H_{p,q}^{m,n}(z)$ has the asymptotic estimate*

(3.17) or (4.25) at infinity.

Remark 6. The power-logarithmic expansions and more complicated results than in (4.7) were indicated in [9, Section 3.7] (see also [8, Section 5.8]) and the particular cases $H_{0,p}^{p,0}(z)$ and $H_{p,p}^{p,0}(z)$ in [7].

Acknowledgement

The authors would like to express their gratitude to the referee for valuable comments.

References

- [1] Braaksma, B.L.G., Asymptotic expansions and analytic continuation for a class of Barnes-integrals, *Compositio Math.* **15**(1964), 239-341.
- [2] Dixon, A.L. and Ferrar, W.L., A class of discontinuous integrals, *Quart. J. Math., Oxford Ser.* **7**(1936), 81-96.
- [3] Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F.G., *Higher Transcendental Functions, Vol. I*, McGraw-Hill, New York, Toronto and London, 1953.
- [4] Fox, C., The G and H functions as symmetrical Fourier kernels., *Trans. Amer. Math. Soc.* **98**(1961), 395-429.
- [5] Kilbas, A.A., Saigo, M. and Shlapakov, S.A., Integral transforms with Fox's H -function in spaces of summable functions, *Integral Transform. Spec. Funct.* **1**(1993), 87-103.
- [6] Marichev, O.I., *Handbook of Integral Transforms of Higher Transcendental Function. Theory and Algorithmic Tables*, Wiley (Ellis Horwood), New York, Brisbane, Chichester and Toronto, 1982.
- [7] Mathai, A.M., An expansion of Meijer's G -function in the logarithmic case with applications, *Math. Nachr.* **48**(1971), 129-139.
- [8] Mathai, A.M., A few results on the exact distributions of certain multivariate statistics. II, *Multivariate Statistical Inference (Proc. Res. Sem. Dalhouse Univ., Halifax, N.S., 1972)*, 169-181, North-Holland, Amsterdam and New York, 1973.
- [9] Mathai, A.M. and Saxena, R.K., *The H -Function with Applications in Statistics and other Disciplines*, Wiley (Halsted P.), New York, London, Sydney and Toronto, 1978.
- [10] Mellin, H.J., Abriß einer einheitlichen Theorie der Gamma- und der hypergeometrischen Funktionen, *Math. Ann.* **68**(1910), 305-337.
- [11] Prudnikov, A.P., Brychkov, Yu.A. and Marichev, O.I., *Integrals and Series, Vol.3, More Special Functions*, Gordon and Breach, New York, Philadelphia, London, Paris, Montreux, Tokyo and Melbourne, 1990.
- [12] Srivastava, H.M., Gupta, K.C. and Goyal, S.P., *The H -Functions of One and Two Variables with Applications*, South Asian Publishers, New Delhi and Madras, 1982.